

Mecánica teórica

pod

Otoño, 2001

Forza generalizada

$$Q_j = \sum \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j}, \quad L(q, \dot{q}, t) = T - V$$

Ec. Lagrange

$$\frac{d}{dt} \left[\frac{\partial T}{\partial \dot{q}_j} \right] - \frac{\partial T}{\partial q_j} = Q_j$$

$$\frac{d}{dt} \left[\frac{\partial L}{\partial \dot{q}_j} \right] - \frac{\partial L}{\partial q_j} = 0$$

Disipativos

$$\frac{d}{dt} \left[\frac{\partial L}{\partial \dot{q}_j} \right] - \frac{\partial L}{\partial q_j} + \frac{\partial \mathcal{F}}{\partial \dot{q}_j} = 0$$

$$\mathcal{F} = \frac{1}{2} \sum (k_x v_{x,i}^2 + k_y v_{y,i}^2 + k_z v_{z,i}^2) = \frac{1}{2} \frac{dW_f}{dt}$$

Potencial general

$$Q_j = -\frac{\partial U}{\partial q_j} + \frac{d}{dt} \left[\frac{\partial U}{\partial \dot{q}_j} \right]$$

Acción $S = \int_{t_1}^{t_2} L(q, \dot{q}, y) dt$, ppi. Hamilton $\delta S = 0$
Momento

$$p_i = \frac{\partial L}{\partial \dot{q}_i}, \quad p_i = \text{cte} \leftrightarrow L \neq L(q_i)$$

f. energía

$$h(q, \dot{q}, t) = \sum_j \dot{q}_j \frac{\partial L}{\partial \dot{q}_j}, \quad \frac{dh}{dt} = -\frac{\partial L}{\partial t}$$

Si $L = L_1 + L_2 + L_3, \rightarrow V \neq V(\dot{q}) \Rightarrow h = E$

Formulismo de Hamilton

$$H(p, q, t) = \dot{q}_i p_i - L(q, \dot{q}, t)$$

$$\dot{q}_i = \frac{\partial H}{\partial p_i}, \quad \dot{p}_i = -\frac{\partial H}{\partial q_i}$$

$$\frac{dH}{dt} = \frac{\partial H}{\partial t} = -\frac{\partial L}{\partial t}, \quad p_i = \frac{\partial L}{\partial \dot{q}_i}$$

$$\begin{aligned} & \text{si } L = L_0 + L_1 + L_2, \quad L_2 = T \quad (q_i \neq q_i(t)) \\ & \rightarrow L_0 = -V \neq -V(\dot{q}), \rightarrow H = L_2 - L_0 = E \end{aligned}$$

$$\begin{aligned} L &= L_0(q, t) + \dot{q}^t \mathbf{a} + \frac{1}{2} \dot{q}^t T \dot{q} \\ H(q, p, t) &= \frac{1}{2} (p^t - \mathbf{a}^t) T^{-1} (p - \mathbf{a}) - L_0(q, t) \end{aligned}$$

$$\delta I = \delta \int_{t_1}^{t_2} L dt = \delta \int_{t_1}^{t_2} (\dot{q}_i p_i - H) dt = 0$$

$$\Delta \int_{t_1}^{t_2} L dt = \int_{t_1 + \Delta t_1}^{t_2 + \Delta t_2} L(\alpha) dt - \int_{t_1}^{t_2} L(0) dt$$

Mínima acción

$$\Delta \int_{t_1}^{t_2} p_i \dot{q}_i dt = 0$$

Transformaciones canónicas. $K = H + \frac{\partial F}{\partial t}$

$$p_i \dot{q}_i - H(q, p, t) = P_i Q_i - K(Q, P, t) + \frac{d}{dt} F(p, P, q, Q, t)$$

$$1. \quad F = F_1(q, Q, t): \quad p_i = \frac{\partial F_1}{\partial q_i}, \quad P = -\frac{\partial F_1}{\partial Q_i}$$

$$\frac{\partial p_i}{\partial Q_k} = \frac{\partial^2 F_1}{\partial Q_k \partial Q_i} = -\frac{\partial P_k}{\partial q_i}$$

$$2. \quad F = F_2(q, P, t) - Q_i P_i. \quad p_i = \frac{\partial F_2}{\partial q_i}, \quad Q_i = \frac{\partial F_2}{\partial P_i}$$

$$\frac{\partial p_i}{\partial P_k} = \frac{\partial^2 F_2}{\partial P_k \partial q_i} = \frac{\partial Q_k}{\partial q_i}$$

$$3. \quad F = q_i p_i + F_3(q, Q, t). \quad q_i = -\frac{\partial F_3}{\partial q_i}, \quad P_i = -\frac{\partial F_3}{\partial Q_i}$$

$$\frac{\partial q_i}{\partial Q_k} = -\frac{\partial^2 F_3}{\partial Q_k \partial p_i} = \frac{\partial P_k}{\partial q_i}$$

$$4. \frac{F}{-\frac{\partial F_4}{\partial p_i}}, \quad Q_i = \frac{\partial F_4}{\partial P_i}$$

$$\frac{\partial q_i}{\partial P_k} = \frac{\partial^2 F_4}{\partial P_k \partial p_i} = -\frac{\partial Q_k}{\partial q_i}$$

$$q_i = \vec{F} = m\vec{a}_f = m\ddot{\vec{R}} + m\vec{a}_r + m\vec{\omega} \times (\vec{\omega} \times \vec{r}) + 2m\vec{\omega} \times \vec{v}_r$$

$$\text{si } \ddot{\vec{R}} = 0 \rightarrow \vec{F}_{\text{ef}} = m\vec{a}_r$$

$$\vec{F}_{\text{ef}} = m\vec{a}_f - m\vec{\omega} \times (\vec{\omega} \times \vec{r}) - 2m\vec{\omega} \times \vec{v}_r$$

Condiciones

$$\begin{aligned} \{Q_i, Q_j\} &= \{q_i, q_j\} = \{P_i, P_j\} = \{p_i, p_j\} = 0 \\ \{P_i, Q_j\} &= \{p_i, q_j\} = \delta_{ij} \end{aligned}$$

Liouville

$$M_{\alpha,\beta} = \frac{\partial x_\alpha}{\partial y_\beta}, \quad M^t J M = J$$

$$\phi_{t,s}(x) = (\varphi_{t,s}^1(x), \dots, \varphi_{t,s}^{2n}(x)), \quad V_s = V_t$$

Paréntesis de Poisson

$$\{f, g\} = \sum_i^n \left(\frac{\partial f}{\partial p_i} \frac{\partial g}{\partial q_i} - \frac{\partial f}{\partial q_i} \frac{\partial g}{\partial p_i} \right)$$

$$\frac{dg}{dt} = \frac{\partial g}{\partial t} + \{H, g\}, \quad \{f \circ \psi, g \circ \psi\} = \{f, g\} \circ \psi$$

Jacobi $\{u, \{v, w\}\} + \{v, \{w, u\}\} + \{w, \{u, v\}\} = 0$

Eq. Hamilton $\dot{q}_k = \{H, q_k\}, \quad \dot{p}_k = \{H, p_k\}$

Hamilton-Jacobi $K = H + \frac{\partial F}{\partial t} = 0, \quad S = F_2$

$$H \left(q_1, \dots, q_n; \frac{\partial S}{\partial q_1}, \dots, \frac{\partial S}{\partial q_n}; t \right) + \frac{\partial S}{\partial t} = 0$$

$$p_i = \frac{\partial S}{\partial q_i} = \alpha_i, \quad Q_i = \frac{\partial S}{\partial P_i} = \frac{\partial S}{\partial \alpha_i} = \beta_i$$

$$S(q, \alpha, t) = W(q, \alpha) - \alpha_1 t, \quad W(q, \alpha) = \sum_i W_i(q_i, \alpha)$$

si $H \neq H(q_j) \rightarrow p_j = P_j = \gamma, \quad W_j = \gamma q_1$

Variables acción-angulo $H(q, p) = \alpha_1 = E$

Momento $J = \oint p dq$

$$Q = \frac{\partial S(q, \alpha, t)}{\partial \alpha_1}, \quad \bar{\omega} = \frac{\partial W(q, J)}{\partial J}$$

$$\dot{\bar{\omega}} = \frac{\partial H(J)}{\partial J} = \nu(J), \quad \bar{\omega} = \int \nu(J) dt = \nu(J)t + \beta$$

S. no inercial

$$\frac{d}{dt} \Big|_f = \frac{d}{dt} \Big|_g + \vec{\omega} \times, \quad \vec{v}_f = \vec{V} + \vec{v}_r + \omega \times \vec{r}$$